

Fennerty Gedge Induction Algorithm Flow Chart

Initiation regimen for RAPID loading (for patients with acute venous thromboembolic disease, e.g. PE / DVT):

Day	INR	Warfarin dose (mg)
1	<1.4	Risk factors* = 5mg No risk factors = 10mg
	≥ 1.4	Seek advice
2	< 1.8	5
	1.8 – 2.0	1
	> 2.0	0
3	< 2.0	5
	2.0 – 2.5	4
	2.6 – 2.9	3
	3.0 – 3.2	2
	3.3 – 3.5	1
	> 3.5	0
4	< 1.4	10
	1.4 – 1.5	7
	1.6 – 1.7	6
	1.8 – 1.9	5
	2.0 – 2.3	4
	2.4 – 3.0	3
	3.1 – 3.2	2
	3.3 – 3.5	1
	> 3.5	0

INRstar takes account of the following *risk factors

- \cdot age > 60 years
- · body weight <50kg
- · liver disease
- · cardiac failure
- · serum albumin <35g/L
- · known bleeding risk
- taking drugs that enhance the effect of anticoagulants
- previously anticoagulated and maintenance dose <2mg daily

a Lumira Dx Care Solutions product

1 North Crofty • Tolvaddon Energy Park • Camborne • Cornwall • TR14 0HX • UK T +44(01209) 710999 F 0845 2412406 E info@INRstar.com www.INRstar.com







